



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

THE NON-REGULAR TRANSITIVE SUBSTITUTION GROUPS WHOSE ORDER IS THE CUBE OF ANY PRIME NUMBER.

By DR. G. A. MILLER, Paris, France.

1. The regular groups of order p^3 (p being any prime number) were published almost simultaneously by Young,* Cole and Glover,† and Hölder.‡ At a considerably earlier date Cayley had published these groups for the special case when $p = 2$.|| It is the object of this paper to determine the remaining transitive groups of this order.

We shall find that there are always two such groups when p is odd. For the special case when p is even it is well known that there is only one.§ Since there are five regular groups of the given order, for every value of p , these facts prove that *the number of the transitive groups of order p^3 is 6 or 7 as p is even or odd.*

The degree, n , ($n < p^3$) of a non-regular transitive group of order p^3 must evidently satisfy the following conditions :

$$p^3 = an, \quad n! = \beta p^3. \quad (a, \beta = \text{positive integers})$$

Hence $n = p^2$. Each one of these groups contains a substitution of order p which is commutative to all its substitutions.¶ Since the cycles of this substitution may be regarded as systems of non-primitivity we see that all these groups are non-primitive.**

The degree of the transitive group which corresponds to the permutations of the systems of non-primitivity when one of these groups is transformed by itself is p . Since the order of this group must be a sub-multiple of p^3 as well as of $p!$ the group must be regular. Hence every non-regular transitive group of order p^3 contains a subgroup of order p^2 which does not interchange the systems of non-primitivity.

The problem of constructing all the possible non-regular transitive groups of order p^3 may therefore be divided into the following two parts: 1) The construction of the intransitive groups of order p^2 which may be used as sub-

* American Journal of Mathematics, XV, p. 133.

† Ibid., p. 196.

‡ Mathematische Annalen, XLIII, p. 371.

|| Philosophical Magazine, XVIII, p. 34; cf. ibid., VII, pp. 40, 408.

§ Serret, Liouville's Journal, 1850, p. 52.

¶ Sylow, Mathematische Annalen, V, 588.

** The transitive groups of order p^a are always non-primitive when $a > 1$.

groups, and 2) The construction of the transitive groups of order p^3 which contain these subgroups. We shall consider these parts separately.

2. *The intransitive groups of order p^2 which may be used as subgroups.*

We shall represent the systems of intransitivity of such a group (H) by

$$A_1, A_2, A_3, \dots, A_p.$$

If one of the substitutions (S) of H would not contain the two systems A_α and A_β ($\alpha, \beta \geq p$) there would be a transform of S with respect to some substitution in the group of order p^3 which would not contain A_α . Since this transform would contain at least one system (A_γ) which is not found in S , it and S would generate a group of order p^2 whose degree would be less than p^2 . All the substitutions of H (excluding identity) must therefore contain either all the systems of intransitivity or all with the exception of one.

The average number of elements in the substitutions of H is $p(p-1)$.* We have just proved that this is also the smallest number of elements that can occur in any substitution of H with the exception of identity. Hence H contains $p-1$ substitutions of degree p^2 , $p(p-1)$ of degree $p(p-1)$, and identity.

If we use one of these substitutions (S_1) of degree p^2 as the first generating substitution of H , the cycles of the p th order in a second generating substitution must be the p different powers of the corresponding cycles in S_1 . Hence there is one and only one H for each value of p . As this H has been completely determined it remains only to consider the construction of

3. *The transitive groups of order p^3 which contain H .*

If we multiply H by a substitution (S_2) which only interchanges its systems according to a cycle of order p we obtain a group (G_1) which contains only substitutions of the p th order and identity† If we multiply H by sS_2 (s being any cycle of S_1) we obtain another group (G_2) which contains $(p+1)$ $(p-1)$ substitutions of order p and $p^2(p-1)$ of order p^2 . It remains to prove that the other groups (G_ϵ ; $\epsilon = 3, 4, \dots$) are not distinct from these two.

Every G is commutative to S_1 . We therefore obtain each one of the possible G 's once and only once if we multiply H by

$$s_1^{a_1} s_2^{a_2} s_3^{a_3} \dots s_p^{a_p} S_2, \quad (a_1, a_2, a_3, \dots, a_p = 1, 2, 3, \dots, p)$$

where $s_1, s_2, s_3, \dots, s_p$ are the cycles of S_1 , in order, and $p' = p-2$.

Since

$$(s_1^{x_1} s_2^{x_2} \dots s_p^{x_p})^{-1} S_2 s_1^{x_1} s_2^{x_2} \dots s_p^{x_p} = s_1^{x_2-x_1} s_2^{x_3-x_2} \dots s_p^{x_1-x_p} S_2,$$

* Frobenius, Crelle's Journal, CI, p. 287.

† The only exception occurs when $p=2$. For this value of p G_1 and G_2 are the same group.

and the system of indeterminate equations

$$x_2 - x_1 = a_1, \quad x_3 - x_2 = a_2, \quad \dots, \quad x_1 - x_p = a_p$$

can always be solved when

$$a_1 + a_2 + a_3 + \dots + a_p = 0,$$

all of these groups must be conjugate to

$$H_{s_1^{a_1} S_2} \quad (a_1 = 1, 2, \dots, p)$$

with respect to substitutions of the form

$$s_1^{a_1} s_2^{a_2} \dots s_p^{a_p}.$$

When $a_1 = 1$, $H_{s_1^{a_1} S_2} = G_2$; when $a_1 = p$, $H_{s_1^{a_1} S_2} = G_1$; for the other values of a_1 , $H_{s_1^{a_1} S_2}$ is evidently conjugate to G_2 with respect to the substitutions which transform S_1 into its different powers without interchanging its cycles. Hence G_1 and G_2 are the only non-regular groups of order p^3 .

APRIL, 1896.